



**DIAGNOSTIC AND INTERVENTIONAL ENDOSCOPY • ERCP • CAPSULE ENDOSCOPY • ACNES**

DR BERNARD CHIN MBBS (Adelaide) FRACP (Australia) Gastroenterologist  
DR JOHN OMBIGA MBBS FRACP (Australia) Gastroenterologist  
DR ANDREW PALMER MBChB (Aberdeen) BSc (Hons) MRCP (UK) FRACP (Aus) Gastroenterologist  
DR MONTRI GURURATSAKUL MD, PhD (Australia), FRACP (Australia) Gastroenterologist

## Medication List

This is a list of the medications that are required to be stopped prior to your upcoming procedure.

Medication	When to STOP
<b>Blood Thinners</b>	
Aspirin & Cartia	Morning of procedure
Apixaban (Eliquis)	2 days prior
Dabigatran (Pradaxa)	2 days prior
Xarelto (Rivaroxaban)	2 days prior
Asasantin	5 days prior
Clopidogrel	5 days prior
Effient (Prasugrel)	5 days prior
Iscover	5 days prior
Plavix	5 days prior
Ticagrelor (Brilinta)	5 days prior
Warfarin (Coumadin/Marevan)	5 days prior
<b>* DO NOT STOP any blood thinners if patient has had a cardiac stent within last 12 months until Gastroenterologist review during prescope appointment</b>	
<b>Others</b>	
Iron tablets	7 days prior
Somac	Morning of procedure
Fish Oil	No need to stop

### For diabetic patients:

Medication	When to STOP
<b>Type I Diabetics</b>	Reduce insulin to 1/3 of regular dose 24 hours prior
<b>Type II Diabetics</b>	Ozempic (Semaglutide, GLp-1)
	7 days prior
	Anti-Hyperglycemic: Dapagliflozin (Forxiga, Qtern, Xigduo XR), Empagliflozin (Jardiance, Glyxambi, Jardimet), Ertugliflozin (Steglatro, Steglujan, Segluromet)
	3 days prior
	Gliptins (Sitagliptin and Vildagliptin)
	24 hours prior
	<b>Any other oral diabetic medications</b> (Byetta, Metformin/Diabex, Janumet, Victoza, Glimepiride) Stop insulin 24 hours prior
	24 hours prior